

Waa Maxay Gelinta Tuubada Gastrojejunostomy (GJ)?

Tubooyinka Mindhiciirka (GJ) waxaa la galiyaa sida caadiga ah iyadoo la isticmaalayo nidaamka loo yaqaan laparoscopy. Laparoscopy waa farsamo lagu sameeyo kamarad u oggolaanaysa bixiyaha inuu arko gudaha ubucda ilmaha. kamarada waxaa la galinayaa meel yar oo la jeexo xudunta hoosteeda. Digi ama tuubo qaabka badhanka ah (GJ-tube) ayaa la galin karaa.

Sida Galinta Tubada Mindhiciirka(GJ) u Caawiso

Ujeedada tuubada GJ waa neefin caloosha cunugeena si ay hawada iyo dheecanka uga saarto. Sidoo kale waxay siisaa wado kale oo cuno ka qaadan karo.



Avanos Mickey GJ



AMT G-JET



Percutaneous Endoscopic Gastrostomy (PEG)

Inta aad isbitaalka joogto

Filo 1 ilaa 2 habeen cisbitaalka inaad joogto. Waxaad ku tababaran doontaa daryeelidda iyo isticmaalka GJ tube si aad u maamusho cuntooyinka iyo daawooyinka inta lagu jiro isbitaal jifidda. Qof cunooyinka ku takhasusay ayaa kaa caawin doona xulashada nooca caanaha ku haboon iyo qiyaasta maalinkasta lasiinayo.

Dhammaan agabka loo baahan yahay si loo daryeelo tuubada iyo quudinta g-tube ka ayaa lagaa siin doonaa shirkadda qalabka caafimaadka (DME) ama shirkadda sahayda guriga ka hor intaadan isbitaalka ka

This information is for educational purposes only. It is not intended to replace the advice of your health care providers. If you have any questions, talk with your doctor or others on your health care team.

If you are a Gillette patient with urgent questions or concerns, please contact Telehealth Nursing at 651-229-3890.

bixin. Waxaa lagu siin doonaa macluumaadka meesha aad kala xiriir karto shirkadda DME, waxaadna awoodi doontaa inaad dalbato sahay dheeri ah mustaqbalka markaad u baahato.

Baxnaaninta iyo soo kabashada

Soo kabashada cunug kastaa way ka duwantahay kan kale, laakiin carruurta intooda badani waxay dib u bilaabaan heerkii dhaqdhaqaaqii caadiga ahaa dhawr maalmood gudahood. Carruurta intooda badan, Tylenol (Acetaminophen) iyo / ama Motrin (Ibuprofen) ayaa ka kaafin doona xanuunka xilliga qalliinka ka dib. Daawooyin ka xoog badan ayaa la isticmaali karaa haddii loo baahdo.

Ku dhaq goobta GJ tubeka maalin kasta saabuun iyo biyo laga bilaabo maalinta qalliinka ka dambeysa. Goobtaas looma baahna in la mariyo, looshan, kareemyo ama boomaato. umana baahna goobta in la dhayo, laakiin hadii aad rabto waxaad isticmaali kartaa faashad. Ha u ogolaanin in faashad qoyan ay sii saarnaato goobta, uga badal sida hadba loogu baahdo.

Ha wareejin tuubada maadaama ay keeni karto inay meesheda ka baxdo. Tuubada xasili. Haddii quudinta ku socoto joogto ama wakhtiyo isku xig xiga, isticmaal aalada kooleyta, sida Flexi-Track si aad ugu xajiso tuubada dheereynta caloosha dusheeda.



Flexi-Track Securement Device



Extension Tube

Ka saar tuubada dheereynta oo dabool tuubada mar kasta oo tuubada aan la isticmaaleynin.

This information is for educational purposes only. It is not intended to replace the advice of your health care providers. If you have any questions, talk with your doctor or others on your health care team.

If you are a Gillette patient with urgent questions or concerns, please contact Telehealth Nursing at 651-229-3890.

ilmahaaga waa in isbuunyo qoyan loogu qubeeyo ama biyaha dusha la iskaga shubo 2 da isbuuc ee ugu horeeya. Qubeyska baafka iyo dabaasha waa caadi 2 isbuuc kadib.

Waxaad u tagi doontaa rugta caafimaadka booqashada raad raaca isbitaal seexinta qalliinka kadib. Jadwalka bedelka tuubada GJ waxaa lagu taliyaa 4 tii ilaa 6 dii biloodba mar in la bedelo.

Haddii aad aragto qandho, guduudasho ama barar ka dhaca goobta GJ Tube ama maqaarka ku xeeran, waxaa laga yaabaa in tan ay calaamad u tahay caabuuq. Wac TeleHealth 651-229-3890.

Waxa loo fiirsado ka dib Markii la Galiyo Tuubada GJ Tube

Granulation Tissue (cad ku dul sameysma): Tani waa dhacdo caadi ah oo G-tube ka weheliso. Cad caafimad qaba ayaa kasoo baxa daloolka si uu daloolka u xidho. Wuxuu u muuqdaa casaan ama guduud, qoyan oo hilib xigeen ah, waxaana ka socda dhacaan qolof dahabi ah. Tani waxaa lagu qiimeyn karaa rugta caafimaadka waxana lagu daweyn karaa Silver Nitrate ama kareem Steroydh ah. Wac si aad ballan u qabsato.



This information is for educational purposes only. It is not intended to replace the advice of your health care providers. If you have any questions, talk with your doctor or others on your health care team.

If you are a Gillette patient with urgent questions or concerns, please contact Telehealth Nursing at 651-229-3890.

Dheecaan: (jaalle, dahabi, bunny, cadaan) oo ka soo baxa tuubada g-tube ka waa la arki karaa waana dheecaanada caloosha ee caadiga ah, waa laga tirtiri karaa ama faashada laga beddeli karaa hadba sidii loogu baahdo.

Daadin: Tani waa dhibaato caadi ah badanaana is xadidda. Daadasho aad u badan micneheedu waxay ka dhigan tahay ilmahaagu mahelayo dareere iyo nafaqo ku filan. Haddii aad aragto daadasho, wac lambarka kalkaalisoooyinka si aad ugala hadasho xeeladaha lagu yareynayo daadinta sida in aad hoos u dhigto quudinta iyo in aad tuubada neefiso. Haddii ay dhacdo daadasho, ka difaac maqaarka xanfah adigoo marinaya zinc oxide (kareemka furuuraca xafaayadda) goobta tuubada, faashadna ku dabool meesha.

Ka saaridda tuubada: g-tube-ka wuu ka soo bixi karaa hadii buufinta haysa ay biyo lumiso ama hadii si kama ah loo soo jiido. Haddii tuubadu ay cusubtahay (in kayar 3 bilood jir) oo ay ka baxdo, buufinka aariyo bixi tuubadana daloolka ku celi. Tuubada ha ku qasbin daloolka. Buufinka dib ha u buufinin. Tuubada meesha ku koolee oo wac lambarka kalkaalisoooyinka, waxayna kuu sheegi doonaan inaad ilmahaaga keentid qeybta gargaarka degdega ah ama xarunta caafimaadka ee Gillette si loogu sameeyo tuubo cusub. Haddii aan tuubo daloolka ku jirin si dhakhso leh ayuu u jufmayaa sidaas darteed dhakhso ku imow qeybta gargaarka degdega ah.

GJ tube kala bedelkiisa: waxaa lagu taliyaa 4 ilaa 6 bilood ama ka hor haddii loo baahdo. Tan waxaa laga yaabaa in qolka qalliinka uu ku sameeyo dhakhtar qalliin ama qeybta Interventional Radiology . Fadlan ballanta u qabso sida uu faray dhakhtarka qalliinka/bixiyaha. Ubedelidda g-tube (caloosha keliya) waxaa laga yaabaa in rugta caafimaadka lagu sameeyo.

Matag: Haddi bukaanka lasoo qaado ama isku dayo inuu matago, dabarka ka fur tuubada caloosha, ama badhanka taabo tuubada dheecaanka ee Bolus exhaust oo aan dabranayn si cadaadiska looga saaro loona oggolaado waxyaabaha caloosha ku jira in ay baxaan. Tani waxaa la yiraahdaa neefin. Haddii bukaanku leeyahay qalliinka hunguriga (Nissen fundoplication), waxaa muhiim u ah in la neefiyo G-tube ka haddii saxasho ama matag ahaan loo soo qaado ay dhacdo. Wac takhtarkaaga haddii saxashada ama soo qaadista matag ahaan joogsan waydo.

This information is for educational purposes only. It is not intended to replace the advice of your health care providers. If you have any questions, talk with your doctor or others on your health care team.

If you are a Gillette patient with urgent questions or concerns, please contact Telehealth Nursing at 651-229-3890.

Xanuun: Haddii bukaanku xanuunsan yahay, quudinta iyo caanaha waxaa laga yaabaa in la bedelo. Weydii takhtarkaaga in ay wanaagsantahay in cunnada laga bedelo.

Xirmid: Haddii tuubadu ay xiranto, wac lambarka kalkaalisoooyinka si aad u hesho tilmaamo. In tuubada xogaa biyo ah la raaciyo inta u dhaxeeyso dawo iyo cunto siinta ayaa ka hor tagta in ay tuubada xiranto.

Quudinta iyada oo loo marinayo Tuubada Caloosha

Kahor quudinta, eeg oo hubi in calooshu madhantahay, haddii dhakhtarka bukaanka uu ku faray in la sameeyo ku dhaji silinge, u oggolow wixii ku jira caloosha dibadda inay usoo qulqulaan, adeegso awooda cuf-jiidka adoo geynaya silingaha meel ka hooseyso heerka caloosha ku taalo, ka dibna silingaha u qabo kor si waxa ku jira caloosha ugu noqdaan.

Si aad u habsatid in tuubadu ay weli si ammaan ah ugu jirto, si tartiib ah dib ugu jiid tuubada ilaa aad dareentid iska caabin yar.

Haddii G-tube ka badhan yahay leh buufin la buufin karo, ha isku dayin inaad si ka duwan siduu yahay ula ekeysiiiso adigoo wax ka badalaya biyahay buufinka. Arag bixiye caafiimaad si sayska wax looga bedelo.

Meelaynta bukaanka

- Udhig qofka bukaanka ah boos raaxo leh, mid toosan ama yara janjeedho. Ilmaha yar waa in la saaro kursiga dhallaanka ama la hayo.
- Waxaa wanaagsan in aan la quudin qofka bukaanka ah, markii si siman uu ujjifo. Haddii loo baahdo u dhig dhinaca midigta.

Quudin Siinta

habka cuf-jiidka / Bolus

This information is for educational purposes only. It is not intended to replace the advice of your health care providers. If you have any questions, talk with your doctor or others on your health care team.

If you are a Gillette patient with urgent questions or concerns, please contact Telehealth Nursing at 651-229-3890.

- Ku dhaji silinge 60ml ah tuubada caloosha.
- Ka buuxi silingaha caanaha quudinta.
- Kor u qaad silingaha iyo tuubada 4 ilaa 4 3/4 inji meel ka sareysa darbiga caloosha.
- Si tartiib ah u quudi, oo qaadanaysa 20 ilaa 30 daqiiqadood.
- Dheecaanka haray ka buuxi ciriiriga ka hor inta uusan madhaneyn, si looga fogaado in hawo uu caloosha galiyo.
- Dareeraha haray ka buuxi silingaha ka hor inta uusan madhanin, si looga hor tago in hawo caloosha gasho.
- HA KA SIININ quudinta cuf-jiidka/bolus tuubada mindhiciirka (GJ Tube)

Habka bamka

- Bacda quudinta ka buuxi qadarka la rabo oo dareere ah.
- Dhareeraha ku shub ilaa dhamaadka tuubada si aad hawada dheeraadka ah uga saarto tubbada.
- Ku xir tuubada tuubada kale ee caloosha ama tuubada mindhiciirka sida lagu faray.
- Sida caadiga ah, quudinta tuubada mindhiciirka gebi ahaanba waxaa la isticmaalaa quudinta bamka mudo dhawr saacadood ha.
- Bilow bamka oo ka dhig heerka ay ku socoto _____ ml saacadiiba..
- Xaqiiji in cunuggaagu uu yahay mid ammaan ah sida ugu macquulsan xiliga quudinta. Waxaad ka hortagi kartaa in si shil ah u saxdo ilmahaaga adigoo tuubada saaraya meel aanu gaari karin, kana fog madaxa, qoorta iyo garbaha inta ay ku xirantahay bacda quudinta.

Quudinta ka dib

- Ku dar _____ ml oo biyo ah bacda quudinta ama silingaha kuna celi nidaamkii quudinta oo kale si loo nadiifiyo tubbada.
- Ka fur silingaha ama tubaada oo dabool saar tuubada caloosha afkeeda.
- Bukaanku waa inuu fadhiistaa 30 daqiiqo kadib quudin kasta.

This information is for educational purposes only. It is not intended to replace the advice of your health care providers. If you have any questions, talk with your doctor or others on your health care team.

If you are a Gillette patient with urgent questions or concerns, please contact Telehealth Nursing at 651-229-3890.

Daawooyinka

- Daawooyinka laga siinayo tuubada caloosha waa inay ahaadaan dareere, ama haddii uu dhakhtarkaada ansixiyey, loo burburiyo oo budo laga dhigo laguna milo dareere. Isku day inaad Ka ilaaliso dawo jadhiidh ah. Waxa ay jufi kartaa G-tube ka.
- Ku sii dawooyinka habka cuf-jiidka adigoo isticmaalaya silinge. Haddii uu yahay G-tube ka qaabka badhanka, isticmaal tuubada bolus extension si aad u siiso dawo la burburiyey. Ha u isticmaalin tuubada xagal toosan leh arrintan.
- Ku falaash 10 ilaa 30 ml oo biyo ah kadib markii aad dawo siiso. Takhtarkaaga ama kalkaalisada ayaa kuu sheegi doona inta biyaha la isticmaalo ayadoo ku saleysan qiyaasta bukaanka iyo baahidiisa dareere.

Kooxda Daryeelkaaga

Bixiyeyaasheena daryeelka caafimaad halkan ayay kuu joogaan markasta xaalad kasta, ee marnaba haka waaban inaad nasoo wacdo.

Kooxda Qalliinka Carruurta ee Gillette Meesha lagala xidhiidhi karo:

- Telehealth 651-229-3890
- Midwest Radiology 651-917-9330
- U of M Interventional Radiology 612-273-6378
- Children's Interventional Radiology 612-874-5399

This information is for educational purposes only. It is not intended to replace the advice of your health care providers. If you have any questions, talk with your doctor or others on your health care team.

If you are a Gillette patient with urgent questions or concerns, please contact Telehealth Nursing at 651-229-3890.